



**Brown Marmorated Stink Bug
Stakeholder Advisory Panel Meeting
USDA/ARS
Appalachian Fruit Research Station
2217 Wiltshire Road
Kearneysville, WV 25430**

December 1, 2015

Submitted by:

Dr. Tracy Leskey

PI, Brown Marmorated Stink Bug Working Group
Research Entomologist
USDA-ARS
Appalachian Fruit Research Station
2217 Wiltshire Road
Kearneysville WV 25430-2771 USA
TEL: 304-725-3451 x329
FAX: 304-728-2340
EMAIL: tracy.leskey@ars.usda.gov

Mr. Guy Moore

Larriland Farms
Fruit and Vegetable Grower
2415 Woodbine Road
Woodbine, Maryland 21797
TEL: 301-854-6110
EMAIL: <http://www.pickyourown.com/contact-form/>
President, Maryland Vegetable Growers Association

Table of Contents

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants.....	3
<i>Introductions and Overview of 2015 Progress</i>	<i>10</i>
<i>Oral Presentation Summaries</i>	<i>10</i>
Objective 1. Voltinism, Dispersal, Landscape and Temporal Risk Factors	10
Objective 1. Nutritional Ecology including diet optimization, salivary gland characterization, gut symbionts and colony procedures.....	10
Objective 1. Commodity Report – commonalities and distinct aspects (injury diagnostics, period of risk, how they invade, level of risk, etc.).....	11
Objective 2. Short-Term Mitigation (Insecticide-Based Management Benefits and Challenges) ...	12
Objective 2. Monitoring Tactics for BMSB	12
Objective 2. Biological Control of BMSB (Native and Classical)	12
Objective 2. Other control tactics (progress on cultural, behavioral and RNAi).....	13
Objective 3. IPM Principles for BMSB management	13
Objective 4. Extension/Outreach efforts at state, regional and national levels.....	14
Objective 4. Economics of BMSB	14
<i>Other Related Projects:</i>	<i>14</i>
OREI	14
SARE.....	15
United Soybean Project	16
Integrating BMSB management into tree fruit IPM	16
<i>Final Discussion</i>	<i>16</i>
<i>Next Steps/Assessments.....</i>	<i>17</i>

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

Name	Affiliation	Title	Full Address
Acebes-Doria, Angel	Virginia Tech	Ph.D. Student	Virginia Tech 216 Price Hall 170 Drillfield Drive Blacksburg, VA 24061-0319
Agnello, Arthur	Cornell NYSAES	Professor	Cornell University/NYSAES 630 West North Street Geneva, NY 14456-1371
Aigner, Ben	Virginia Tech	M.S. Student	Virginia Tech 216 Price Hall 170 Drillfield Drive Blacksburg, VA 24061-0319
Aigner, John	Virginia Tech	Ph.D. Student	Virginia Tech 216 Price Hall 170 Drillfield Drive Blacksburg, VA 24061-0319
Beck, David (Webinar)	Crawford Beck Vineyard, LLC	Vineyard Owner	Crawford Beck Vineyard, LLC 9100 SE Amity Road Amity, OR 97101
Behling, George	Nob Hill Orchards	Tree Fruit Grower & Owner	Nob Hill Orchards 1572 Reunion Corner Road Gerrardstown, WV 25420
Bergh, Chris	Virginia Tech	Professor	Alson H. Smith, Jr. AREC 595 Laurel Grove Road Winchester, VA 22602
Blaauw, Brett	Rutgers University	Post-doctoral Research Associate	Rutgers University 121 Northville Road Bridgeton, NJ 08302

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

Name	Affiliation	Title	Full Address
Brandt, Nathaniel	USDA-ARS-AFRS	Undergraduate Student	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Bush, Hayley	USDA-ARS-AFRS	Undergraduate Student	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Cullum, John	USDA-ARS-AFRS	M.S. Student	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Dieckhoff, Christine	UDEL/USDA-ARS-BIIR	Visiting Scientist	USDA/ARS/BIIRU 501 South Chapel Street Newark, DE 19713
Dively, Galen	University of Maryland	Professor Emeritus and IPM Consultant	University of Maryland 4112 Plant Sciences Building College Park, MD 20742-4454
Futrell, Susan	Red Tomato	Marketing Director	Red Tomato 76 Everett Skinner Road Plainville, MA 02762
Gauen, Kenneth	The Pictsweet Company	Field Representative	The Pictsweet Company PO Box 398 Bridgeville, DE 19933
Gonzales, Chris	Cornell University	Communication Specialist	Cornell University Northeastern IPM Center 169 Helios Circle The Insectary, Blauvelt Lab Ithaca, NY 14853
Green, Tom	IPM Institute of North America	President	IPM Institute of North America 1020 Regent St Madison, WI 53715

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

Name	Affiliation	Title	Full Address
Haas, Tom	Cherry Hill Orchards	Tree Fruit Grower & Owner	Cherry Hill Orchards 2194 New Danville Pike Lancaster, PA 17603
Hamilton, George	Rutgers University	Specialist in Pest Management	Rutgers University 93 Lipman Drive New Brunswick, NJ 08901
Hancock, Torri	USDA-ARS-AFRS	M.S. Student	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Harper, Jayson	Penn State University	Professor	Penn State University 201-D Armsby Building University Park, PA 16802
Hoffman, Scott	Furmano Foods	Field Representative	Furmano Foods 770 Cannery Road Northumberland, PA 17857
Hollabaugh, Brad	Hollabaugh Brothers	Tree Fruit Grower	Hollabaugh Brothers, Inc 545 Carlisle Road Biglerville, PA 17307
Inkley, Douglas	National Wildlife Federation	Senior Scientist	National Wildlife Federation 11100 Wildlife Center Drive Reston, VA 20190
Jones, Sharon	USDA-ARS-AFRS	Technician	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Joy, Donna	USDA-ARS-AFRS	Program Support	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

Name	Affiliation	Title	Full Address
Klimstra, Gene	Agricon Consulting	Independent Consultant	Agricon Consulting 5730 Willow Road Hendersonville, NC 28739
Krawczyk, Grzegorz	Penn State University	Associate Professor	Penn State University/FREC 290 University Drive Biglerville, PA 17307
Kuhar, Thomas	Virginia Tech	Professor	Virginia Tech 170 Drillfield Drive Blacksburg, VA 24061-0319
Leskey, Tracy	USDA-ARS-AFRS	Entomologist	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Liu, Yifen	Northeastern IPM Center	Evaluation Specialist	Cornell University Northeastern IPM Center The Insectary Ithaca, NY 14853
Maccherone, Santo John (Ricky)	Circle M Fruit Farms	Grower	Circle M Fruit Farms 88 Sandy Ridge Rd Mannington Township, NJ 08079
MacKintosh, Bill	MacKintosh Fruit Farm	Fruit Farm Grower & Owner	MacKintosh Fruit Farm 1608 Russell Road Berryville, VA 22611
Martinson, Holly	University of Maryland	Student	University of Maryland 4112 Plant Sciences Building College Park, MD 20742
Moore, Guy	Larriland Farms	Fruit & Vegetable Grower	Larriland Farms 2415 Woodbine Road Woodbine, MD 21797
Morehead, Adam	Virginia Tech	M.S. Student	Virginia Tech 170 Drillfield Drive Blacksburg, VA 24061-0319

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

Name	Affiliation	Title	Full Address
Morrison, Rob	USDA-ARS-AFRS	Post-doctoral Research Associate	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Newkirk, Timothy	Willamette Hazelnut	Grower	Willamette Hazelnut 14975 NE Tangen Road Newberg, Oregon 97132
Nicolosi, Brandon	USDA-ARS-AFRS	Undergraduate Student	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Nielsen, Anne	Rutgers University	Assistant Extension Specialist	Rutgers University 121 Northville Road Bridgeton, NJ 08302
Orr, Mark	Orr's Farm WV	Grower	Orr's Farm Market 682 Orr Drive Martinsburg, WV 25403
Poling, Brittany	USDA-ARS-AFRS	Undergraduate Student	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Rentzel, Kay	National Peach Council	Managing Director	National Peach Council 22 Triplett Court Dillsburg, PA 17019
Rice, Kevin	USDA-ARS-AFRS	Post-doctoral Research Associate	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Rodriguez-Saona, Cesar	Rutgers University	Associate Extension Specialist	Rutgers University P.E. Marucci Center 125A Lake Oswego Road Chatsworth, NJ 08019
Rice, Kevin	USDA-ARS-AFRS	Post-doctoral Research Associate	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

Name	Affiliation	Title	Full Address
Seagraves, Michael (Webinar)	Driscoll's	Entomology Lead	Driscoll's 151 Silliman Road Watsonville, CA 95076
Shearer, Peter	Oregon State University	Professor	OSU Mid-Columbia Agricultural Research & Extension Center 3005 Experiment Station Drive Hood River, OR 97031
Shenot, Rob (Webinar)	Shenot Farms	Fruit & Vegetable Grower	Shenot Farms 201 Peachstone Lane Wexford, PA 15090
Short, Brent	USDA-ARS- AFRS	Researcher	USDA/ARS/AFRS 2217 Wiltshire Road Kearneysville, WV 25430
Showalter, Lee	Rice Fruit Co.	Food Safety Manager/ Grower Services	Rice Fruit Co. 2760 Carlisle Road Gardners, PA 17324
Sisco, Rebecca (Webinar)	Western Region IR-4 Center	Regional Field Coordinator	Western Region IR-4 Center 4218 Meyer Hall NE Corner La Rue & Bioletti Way UC Davis, CA 95616
Taylor, Christopher	University of Maryland	Ph.D. Student	University of Maryland 4112 Plant Sciences Building College Park, MD 20742
Walgenbach, Jim	NC State University	Extension Entomology Specialist	MHCREC 455 Research Drive Mills River, NC 28759
Walton, Vaughn	Oregon State University	Associate Professor	Oregon State University 4109 Ag & Life Sci Building Corvallis, OR 97331

Brown Marmorated Stink Bug Stakeholder Advisory Panel Participants

<u>Name</u>	<u>Affiliation</u>	<u>Title</u>	<u>Full Address</u>
Whalen, Joanne	University of Delaware Cooperative Extension	Extension Specialist State Program Leader, Agriculture and Natural Resources	College of Agriculture & Natural Resources 531 South College Ave. Newark, DE 19716
Wise, John	Michigan State Univ	Associate Professor	Michigan State University CIPS 578 Wilson Road East Lansing, MI 48824

Introductions and Overview of 2015 Progress

Presented by: Tracy Leskey, USDA/ARS/AFRS

Summary:

- Introduction of each person in attendance.
- Information for attendees regarding the contents of program packet provided.
- Presentation on Stakeholder Advisory Panel purposes, objectives, funding, and accomplishments to date.
- Reviewed published manuscripts which supported each objective and reviewed each objectives' progress along with further pending questions
- Presented number of key personnel trained
- Feedback from 2014 SAP meeting reviewed

Oral Presentation Summaries

Objective 1. Voltinism, Dispersal, Landscape and Temporal Risk Factors

Presented by: Anne Nielsen, Rutgers University

Summary:

- Overwintering BMSB emerged over a period of ~2.5 months
- Early, smaller peak of emergence appeared to be primarily associated with a period of warmer temperatures (smaller and/or weak bugs depleted of resources?)
- Later, larger peak associated with temperature and/or change (stabilized?) in photoperiod
- Pheromone traps appeared to reflect onset emergence reasonably well
- Marked bugs assumed to have dispersed from emergence site
- Need to conduct studies in north-south transect with naturally settling adults
- Individual-based model developed to predict voltinism.

Objective 1. Nutritional Ecology including diet optimization, salivary gland characterization, gut symbionts and colony procedures

Presented by: Chris Taylor, University of Maryland and Nik Wiman, Oregon State University

Summary:

- Adults emerging from overwintering exhibit a steady decline in lipids, glycogen, and sugars as they emerged later in the season. This suggests that overwintering for longer periods of time uses up more nutrient reserves.
- Adults that emerged from overwintering in May-June had numerically lower weights and nutrient reserves than their counterparts collected from holly in OR at the same time. This suggests that feeding on host plants may have replenished their reserves.
- Mixed host plant diets proved to be optimal for nymphal survivorship and development
- Nymphs reared on mixed diets and ToH developed faster and they resulted into bigger and heavier adults

- Peach appeared to be the most suitable single host for BMSB development among the host plants tested
- Nutrient levels of adults that developed from nymphs reared on different diets were different across treatments
- Results suggest that *H. halys* optimizes diet by utilizing multiple hosts during its development
- Colony Rearing and Diapause - Summary of findings
 - Optimal temperature and humidity
 - Mixed diet of proteins and carbohydrates
 - Diapause considerations
 - Issues with microsporidian infection

Objective 1. Commodity Report – commonalities and distinct aspects (injury diagnostics, period of risk, how they invade, level of risk, etc.)

Presented by: Chris Bergh, Virginia Polytechnic Institute and State University

Summary:

- Feeding by 2nd & 3rd instar BMSB nymphs on apples and pears did not cause as much injury as feeding by 4th & 5th instars or adults
- BMSB abundance on fruiting ornamentals tracked seasonal availability of fruit. De-fruiting trees caused pronounced reduction in BMSB abundance
- Initial indications that feeding by BMSB on Chardonnay and Pinot Noir grapes tended to make them more vulnerable to SWD infestation
- Damage on Bosc more severe than on d’Anjou pears from same exposure to BMSB just before harvest
- BMSB feeding injury increased ethylene production and respiration rate in Bosc but not d’Anjou pears during cold storage

Discussion Break:

- Photoperiod inclusion in individual-based model – Homestead FL data –populations will not be severe.
- Are minerals important for the development of BMSB – unknown
- If you find microsporidia in your colonies – contact Anne Hajek at Cornell, as these could be used as a biocontrol measure. Field collections showed high 20 to 25 percent parasitism – this may help biocontrol
- More BMSB overwintering – outside overwintering, more bugs, more forests. Has to be overwintering – 35,000 bugs shed at top of mountain
- Pollinator concerns may result in loss of neonics in management program. Neonics – EPA to restrict for applied – application on labels very restrictive. Management plans need to adapt and give growers alternative tools.
- Standardization of traps and lures for each specialty crop, but likely not same across crops.
- Insecticides work – possible resistance. Pyrethroids/neonicotinoids. Baseline susceptibility needs to be tracked – consider baseline – impact of insecticides

Objective 2. Short-Term Mitigation (Insecticide-Based Management Benefits and Challenges)

Presented by: Greg Krawczyk, Penn State University

Summary:

- Most effective insecticides against BMSB
 - Pyrethroids (IRAC Group 3A)
 - Bifenthrin (Brigade)
 - Fenpropathrin (Danitol)
 - Cyfluthrin (Baythroid)
 - λ -cyhalothrin (Warrior)
 - Neonicotinoids (IRAC Group 4A)
 - Dinotefuran (Venom, Scorpion)
 - Thiametoxam (Actara)
 - Clothianidin (Belay)
 - Imidacloprid (Provado, Admire Pro)
 - Acetamiprid (Assail)
 - Other (IRAC Groups 1A, 1B, 2A)
 - Methomyl (carbamate) (Lannate LV and SP)

Objective 2. Monitoring Tactics for BMSB

Presented by: Tracy Leskey, USDA

Summary:

- Identification and commercialization of BMSB aggregation pheromone and pheromone synergist
- Early season attraction documented for BMSB from March 20-April 17, 2012
- Two-component BMSB aggregation pheromone identified - Khrimian et al. 2014
- No significant difference in BMSB responses to varying levels of purity
- Collaborations with commercial companies throughout the project.
- Provided commercial collaborators with samples of BMSB pheromone for formulation and testing.
- Coordinated lure trials in 2014 and 2015 with current commercial formulations.
- Most lures perform equally as well as the experimental standard.
- Traps provisioned with white light sources were most attractive in the field, but blue light sources were most specific for BMSB

Objective 2. Biological Control of BMSB (Native and Classical)

Presented by: Christine Dieckhoff, USDA

Summary:

- *Trissolcus japonicus* is oligophagous - it attacks several Asian pentatomid species
- These populations are adventive – they were not released nor did they escape quarantine!

- DC area populations are genetically similar to populations sampled in Japan and S. Korea
- WA population is genetically similar to populations sampled in S. Korea
- On average, about 40-50% of the eggs we found of other stink bugs were parasitized; i.e. 10-fold the number of eggs parasitized compared to naturally laid BMSB eggs. Most of these (~70%) are Scelionidae – Trissolcus and Telenomus spp.
- Parasitism of sentinel and naturally laid eggs low by native natural enemies throughout season, across years and locations (2006/2007 through 2015)
- Survey of parasitism of non-BMSB naturally laid eggs suggests that native egg parasitoid species are present and active in the same habitats as BMSB
- Species composition of sentinel and naturally laid eggs:
 - dominated by *Anastatus* spp. (Eupelmidae)
 - highly variable between years
 - proportion of undeveloped eggs highly variable between year, partly due to state of egg mass at deployment (frozen vs. fresh)

Objective 2. Other control tactics (progress on cultural, behavioral and RNAi)

Presented by: Rob Morrison, USDA

Summary:

- The major BMSB secretion compound (C13) has no effect on feeding activity.
- E2-decanal (E2) demonstrate dual functions in petri dish assay. At low concentration (>100 µg), it acts as feeding stimulant; while at high concentration (<100 µg), it acts as feeding deterrent.
- A blend of tridecane (C13) and E2-decanal (E2) (1:1 ratio) showed significant feeding deterrent activity.
- Isolongifolenone (En), a mosquito and tick repellent, exhibits strong feeding deterrent activity.
- A blend of C13, E2, and En (1:1:1 ratio) also exhibits strong feeding deterrent activity.
- Although identified feeding deterrents are natural products, they cannot be directly applied on the plants (surface burning). New formulation is ready to be tested in the field this fall.
- The identified feeding deterrents/repellents can be easily commercialized and used for protecting agricultural crops from *H. halys* damage in support of ongoing *H. halys* management programs.

Objective 3. IPM Principles for BMSB management

Presented by: Tom Kuhar, Virginia Polytechnic Institute and State University

Summary:

- BMSB is a border-driven pest and we should focus on border-driven management tactics – border sprays, attract and kill, etc.
- Monitoring tools being developed for specialty crops
- Trap captures reflected relative, local densities of BMSB in various studies.

Threshold = 10 adults per trap resulted in a 40% reduction in insecticide applications with injury at harvest statistically equivalent to blocks treated weekly in apple. Baited traps can be used to guide management decisions in apple orchards.

- Net structures fall into three basic types depending on their use: A completely enclosed orchard, that a tractor can be driven through, has both top and sides. A sturdy structure (posts and wires) is required to support the nets and entryways.

Objective 4. Extension/Outreach efforts at state, regional and national levels.

Presented by: Chris Gonzales, Northeastern IPM Center

Summary:

- 3,500 stakeholders on Center e-mail list
- 375 recipients on BMSB e-mail lists
- 4,600 regional addresses receive print
- Facebook: 267 likes
- Twitter: 1,911 followers / 225 retweets (past year)
- YouTube (over 27,000 views in past year, 52 subscribers)
- Flipboard (81 viewers)

Objective 4. Economics of BMSB

Presented by: Jay Harper, Penn State University

Summary:

- Penn State Logo Change
- More data on the impact of BMSB on the cost of producing apples, peaches, and tomatoes will be evaluated in 2016.
- Will be helping determine the costs and potential benefits of proposed management tactics in 2016 (and beyond).
 - Estimate cost and benefits of proposed management strategies and make available through extension channels.
 - Evaluate potential physical and financial constraints faced by producers in implementing the proposed tactics.
 - Fine tune management recommendations and provide feedback to producers on the status and commercial viability of proposed control strategies.

Other Related Projects:

OREI

Presented by: Anne Nielsen, Rutgers University

Summary:

- Sorghum was generally the most attractive trap crop tested for BMSB

- Sunflower was more attractive earlier in the season with sorghum becoming more attractive in August
- Sunflower is attractive to natural enemies
- Colonization of cash crop was delayed
- Higher damage in peppers occurred under ‘high’ pressure
- Also attractive to native stink bugs
- Nymphs have a strong walking capacity.
- Can disperse 10m in 3 hours
- Nymphs show strong response to the olfactory attractant and traverse large distances to reach source
- Nymphs select host plants
- Based off of phenology
 - Preference for fruiting bodies
 - Identified common odors correlated with attraction
- Egg mass predation is higher in organic systems than conventional
- Most predators are generalists or opportunists
 - Sucking predators, orthopterans
- Can be increased through habitat manipulation
 - Until *T. japonicus* is widespread, focus should be on plants that increase predator community
 - Horsemint (*Monarda* sp) and Coreopsis
 - Insecticides like Entrust decrease NE populations
- Parasitism is increasing
- Is Organic Management Feasible?
 - Yes, *under moderate pressure!*
- Understand hot spots on the farm
 - Key early season host plants
 - Crops that are preferred hosts by all life stages
- Manipulate the habitat surrounding these areas
 - Support natural enemies
 - Trap crop using sunflower and sorghum
 - Re-design trap crop layout
- Under intense BMSB pressure the finest mesh netting provides protection from stink bug injury
- Remove overwintering populations on-farm

SARE Attract and Kill System for Apple

Presented by: Brent Short, USDA

Summary:

- Lower cost lures
 - Competition/Volume/Refinement of production
 - Smaller doses

- Inclusion of Host Plant Volatiles
- Fewer lures
 - Fewer per tree and/or fewer baited trees
- Less frequent spray applications
 - Threshold + Perimeter-Based Management
- Impact on farm-level management
- Long-term reduction of BMSB populations

United Soybean Project

Presented by: Galen Dively, University of Maryland

Summary:

- Soybean yield sampling – drop off/2 min visual count threshold
- Difference in temperance – aspect slope
- Pocket guide – card with lanyard
- Reduced yield – doesn't fill pod – wait until it turns color
- Presented project results and information on the biology and management of BMSB at 13 invited out-of-state crop conferences and professional events and 58 in-state producer meetings
- Six scientific papers on the results of the project are either in the review process or published.
- Produced an extension field guide that summarizes much of what was accomplished in this project. <http://unitedsoybean.org/brown-marmorated-stink-bugs/>

Integrating BMSB management into tree fruit IPM

Presented by: Brett Blaauw, Rutgers University

Summary:

- IPM-CPR reduces insecticide use by up to 75%
- Pest control and fruit damage at levels equal to current management recommendations
- Promising data in apples as well
- May reduce negative impact on natural enemies
 - Important for secondary pests?

Final Discussion

- 25 percent injury – late start in applying – near forests – using pheromone trap – increased 4 traps 16 sampling sites – 8-10 percent really late season
- go for additional grant
- Bring everyone together after field season
- End Aug 2016 – go on by webinar
- Helpful counts – move ahead – develop threshold – trap and lure standard

- Project – reduce species – invasive species – stronger controls – will talk to you about it – agenda again with less insecticide
- Working Group meeting will continue – growers invited and can still come
- Very impressed with collaborator communication – real gift – appreciate appointment and to see what is going on with BMSB and as a Stakeholder Advisory Panel member

Next Steps/Assessments

- Wrapping up – Final grant ends August 31, 2016
- Hope to use the input from each of you for grading objectives/priorities
- Three pieces –
 - SCRI Stakeholder Review - Standard Feedback Form
 - BMSB NEIPM Priorities
 - BMSB Western Priorities
- Smithsonian – Genome *Japonicus* Grant submitted
- OECD funding in Europe?
- OREI/OREF/SARE
- Stop Pest/HUD/NPM
- Private Donors
- Gates Foundation
- Specialty Crop Block grants
- Research and extension personnel were given thanks for being such hard workers
- Nice Job! Team effort